

Supergravity from Bosonic String

B. B. Deo

Department of Physics, Utkal University, Bhubaneswar, India

(Dated: February 1, 2008)

Deser-Zumino [1] N=1 supergravity action for the graviton and gravitino pair in four dimensions is deduced from the Nambu-Goto action [2, 3] of the 26 dimensional bosonic string theory using Principle of Equivalence.

PACS numbers: 04.65.+e, 12.60.J

Keywords: supergravity, supersymmetry, superstring

Not long ago, Casher, Englert, Nicolai and Taormina [4, 5] showed that consistent superstrings can be the solutions of D=26 bosonic string theory and the later appears as the fundamental string theory. This has also been demonstrated by me and in collaboration with L. Maharana in recent references [6] and [7, 8, 9]. Details of construction of consistent superstring are given there. None of the existing four dimensional string theories contain construction of gravitino and graviton ground states from string theory excitation quanta leading to supergravity. As far as I know, this is the first attempt in this direction. I repeat some of them so that the preprints can be skipped without missing the deductions of the main theme which is of immediate information to string theorists. The physical open bosonic string theory with the coordinates X^μ in the world sheet (σ, τ) is given by the action

$$S = -\frac{1}{2\pi} \int d^2\sigma \partial^\alpha X^\mu(\sigma, \tau) \partial_\alpha X_\mu(\sigma, \tau), \quad \mu = 0, 1, 2, \dots, 25. \quad (1)$$

It is well known that we can introduce 44 Majorana fermions ϕ_λ as proven basically by Mandelstam [10] for (1+1) quantum field theory, which are scalars in the world sheet and four bosonic coordinates and the action

$$S = -\frac{1}{2\pi} \int d^2\sigma \left[\partial^\alpha X^\mu(\sigma, \tau) \partial_\alpha X_\mu(\sigma, \tau) - i \sum_{j=1}^{44} \bar{\phi}^j \rho^\alpha \partial_\alpha \phi_j \right], \quad \mu = 0, 1, 2, 3. \quad (2)$$

where we have taken

$$\partial_\alpha = (\partial_\sigma, \partial_\tau), \quad \rho^0 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \rho^1 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \bar{\phi} = \phi^\dagger \rho^0 \quad (3)$$

In the Chapter-6.1.2 in [11], it has also been proved and stated that two Majorana fermions in 1+1 dimensions are equivalent to one real boson in the infinite volume limit. It is also true in anomaly free string theory on a finite interval or a circle even one like in Veneziano model. Observing that there are also Majorana Lorentz fermions ψ^μ in the bosonic representation of SO(3,1), we can further rewrite equation (2) as [6]

$$S = -\frac{1}{2\pi} \int d^2\sigma \left[\partial^\alpha X^\mu(\sigma, \tau) \partial_\alpha X_\mu(\sigma, \tau) - i \sum_{j=1}^{11} \bar{\psi}^{\mu,j} \rho^\alpha \partial_\alpha \psi_{\mu,j} \right] \quad (4)$$

presuming this to be anomaly free. This has been proved in reference [6]. The upper index refer to row and the lower index to a column. Searching for a Ψ^μ , to copartner to X^μ , one has to further divide the elevens into two groups. There is a group of six, $j=1,2,\dots,6$, where the negative and positive frequency parts are $\psi^{\mu,j} = \psi^{(+)\mu,j} + \psi^{(-)\mu,j}$ and another group of five, $k=7,8,\dots,11$ with $\phi^{\mu,k} = \phi^{+\mu,k} - \phi^{-\mu,k}$. The negative sign is absorbed in creation operators as allowed by the freedom of phase of Majorana fermions. I introduce matrices (e^j, e^k) which are rows of ten zeroes and only one 1 in the j^{th} or k^{th} place. $e^j e_j = 6$ and $e^k e_k = 5$. With these the world sheet action,

$$S = -\frac{1}{2\pi} \int d^2\sigma \left[\partial^\alpha X^\mu(\sigma, \tau) \partial_\alpha X_\mu(\sigma, \tau) - i \bar{\psi}^{\mu,j} \rho^\alpha \partial_\alpha \psi_{\mu,j} + i \bar{\phi}^{\mu,k} \rho^\alpha \partial_\alpha \phi_{\mu,k} \right] \quad (5)$$

is invariant under supersymmetric transformations,

$$\delta X^\mu = \bar{\epsilon} (e^j \psi_j^\mu - e^k \phi_k^\mu), \quad (6)$$

$$\delta \psi^{\mu,j} = -i \epsilon e^j \rho^\alpha \partial_\alpha X^\mu, \quad (7)$$

$$\text{and} \quad \delta \phi^{\mu,k} = -i \epsilon e^k \rho^\alpha \partial_\alpha X^\mu. \quad (8)$$

ϵ is a constant anticommuting spinor.

In action (5), there are 4 bosons but eleven fermions which are Lorentz vectors of SO(3,1) which is possible only in 1+1 dimensions [11]. This will show up in two supersymmetric successive transformations. In our case, in particular, the transformation leads to spatial translation with the coefficient $a^\alpha = 2i\epsilon^1 \rho^\alpha \epsilon_2$ but constrained by

$$\psi_j^\mu = e_j \Psi^\mu, \quad \phi_k^\mu = e_k \Psi^\mu \quad (9)$$

which defines the nature of e_j and e_k and also provides the much needed linear combination for the fermion SUSY partner of X^μ

$$\Psi^\mu = e^j \psi_j^\mu - e^k \phi_k^\mu. \quad (10)$$

There is no need for additional auxilliary field. The action (5) reduces to

$$S = -\frac{1}{2\pi} \int d^2\sigma [\partial^\alpha X^\mu(\sigma, \tau) \partial_\alpha X_\mu(\sigma, \tau) - i\bar{\Psi}^\mu \rho^\alpha \partial_\alpha \Psi_\mu] \quad (11)$$

Using equations (6) to (8) and equation (11) itself, it is easy to verify

$$\delta X^\mu = \bar{\epsilon} \Psi^\mu, \quad \delta \Psi^\mu = -ie\rho^\alpha \partial_\alpha X^\mu \quad (12)$$

$$[\delta_1, \delta_2] X^\mu = a^\alpha \partial_\alpha X^\mu, \quad [\delta_1, \delta_2] \Psi^{\mu,j} = a^\alpha \partial_\alpha \Psi^{\mu,j}. \quad (13)$$

The actions given in equations(5) and (13) are equivalent. Ψ^μ , while in j^{th} or k^{th} site emits and absorbs quanta by quantising $\psi^{\mu,j}$ or $\phi^{\mu,k}$ according to the equation (5) which should be unitary, free of ghosts and nomalies. It should be free of tachyons and modular invariant. These aspects have been studied in detail in references [9] and [7], where, also currents and energy momentum tensors have been given in light cone basis as deduced from local 2d-global symmetric extension of the above action. We shall not need these details here.

The action in (11) yields the only nonvanishing equal time commutator and anticommutators [11] as

$$[\partial_\pm X^\mu(\sigma, \tau), \partial_\pm X^\nu(\sigma', \tau)] = \pm \frac{\pi}{2} \eta^{\mu\nu} \delta'(\sigma - \sigma') \quad (14)$$

$$\{\psi_A^\mu(\sigma, \tau), \psi_B^\nu(\sigma', \tau)\} = \pi \eta^{\mu\nu} \delta(\sigma - \sigma') \delta_{AB} \quad (15)$$

$$\{\phi_A^\mu(\sigma, \tau), \phi_B^\nu(\sigma', \tau)\} = -\pi \eta^{\mu\nu} \delta(\sigma - \sigma') \delta_{AB} \quad (16)$$

$$\{\Psi_A^\mu(\sigma, \tau), \Psi_B^\nu(\sigma', \tau)\} = \pi \eta^{\mu\nu} \delta(\sigma - \sigma') \delta_{AB} \quad (17)$$

On a light cone basis, the Noether supercurrents and energy momentum tensors are

$$J_+ = \partial_+ X_\mu \Psi_+^\mu \quad (18)$$

$$J_- = \partial_- X_\mu \Psi_-^\mu \quad (19)$$

$$T_{++} = \partial_+ X^\mu \partial_+ X_\mu + \frac{i}{2} \bar{\psi}_+^{\mu,j} \partial_+ \psi_{+\mu,j} - \frac{i}{2} \bar{\phi}_+^{\mu,k} \partial_+ \phi_{+\mu,k} \quad (20)$$

$$T_{--} = \partial_- X^\mu \partial_- X_\mu + \frac{i}{2} \bar{\psi}_-^{\mu,j} \partial_- \psi_{-\mu,j} - \frac{i}{2} \bar{\phi}_-^{\mu,k} \partial_- \phi_{-\mu,k} \quad (21)$$

It follows from the variation of the local 2-d supersymmetric action that the above equations (18) to (21) vanish. This enables one to construct physical states without ghosts in the Fock space. One quantises the action (5) as [6]

$$X^\mu(\sigma, \tau) = x^\mu + p^\mu \tau + i \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu e^{-in\tau} \cos(n\sigma) \quad (22)$$

$$\text{or} \quad \partial_\pm X^\mu = \frac{1}{2} \sum_{-\infty}^{+\infty} \alpha_n^\mu e^{-in(\tau \pm \sigma)} \quad (23)$$

$$\text{and} \quad [\alpha_m^\mu, \alpha_n^\nu] = m \delta_{m+n} \eta^{\mu\nu} \quad (24)$$

With Neveu-Schwartz(NS) and Ramond (R) [13, 14] boundary conditions

$$\Psi_\pm^\mu = \frac{1}{\sqrt{2}} \sum_{r \in Z + \frac{1}{2}} B_r^\mu e^{-ir(\tau \pm \sigma)}, \quad \text{with} \quad b_r^{\mu,j} = e^j B_r^\mu \quad \text{and} \quad b'^{\mu,k} = e^k B_r^\mu \quad \text{NS} \quad (25)$$

$$= \frac{1}{\sqrt{2}} \sum_m D_m^\mu e^{-im(\tau \pm \sigma)}, \quad \text{with} \quad d_m^{\mu,j} = e^j B_m^\mu, \quad \text{and} \quad d'^{\mu,k} = e^k B_m^\mu \quad \text{R} \quad (26)$$

and the anticommutation relations are

$$\{B_r^\mu, B_s^\nu\} = \eta^{\mu\nu} \delta_{r+s}, \quad \{D_m^\mu, D_n^\nu\} = \eta^{\mu\nu} \delta_{m+n}. \quad (27)$$

To construct admissible states, it is essential to give the Super Virasoro generators and their algebra in notations of reference [13, 14]

$$L_m = \frac{1}{\pi} \int_{-\pi}^{\pi} d\sigma e^{im\sigma} T_{++} \quad (28)$$

$$= \frac{1}{2} \sum_{-\infty}^{\infty} : \alpha_{-n} \cdot \alpha_{m+n} : + \frac{1}{2} \sum_{r \in z + \frac{1}{2}} (r + \frac{1}{2}m) : (b_{-r} \cdot b_{m+r} - b'_{-r} \cdot b'_{m+r}) : \quad \text{NS} \quad (29)$$

$$= \frac{1}{2} \sum_{-\infty}^{\infty} : \alpha_{-n} \cdot \alpha_{m+n} : + \frac{1}{2} \sum_{n=-\infty}^{\infty} (n + \frac{1}{2}m) : (d_{-n} \cdot d_{m+n} - d'_{-n} \cdot d'_{m+n}) : \quad \text{R} \quad (30)$$

$$G_r = \frac{\sqrt{2}}{\pi} \int_{-\pi}^{\pi} d\sigma e^{ir\sigma} J_+ = \sum_{n=-\infty}^{\infty} \alpha_{-n} \cdot (e^j b_{n+r,j} - e^k b'_{n+r,k}), \quad \text{NS} \quad (31)$$

$$F_m = \sum_{-\infty}^{\infty} \alpha_{-n} \cdot (e^j b_{n+r,j} - e^k b'_{n+r,k}). \quad \text{R} \quad (32)$$

The Super Virasoro algebra is

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{C}{12}(m^3 - m)\delta_{m,-n}, \quad (33)$$

$$[L_m, G_r] = (\frac{1}{2}m - r)G_{m+r}, \quad \text{NS} \quad (34)$$

$$\{G_r, G_s\} = 2L_{s+r} + \frac{C}{3}(r^2 - \frac{1}{4})\delta_{r,-s}, \quad (35)$$

$$[L_m, F_n] = (\frac{1}{2}m - n)F_{m+n}, \quad \text{R} \quad (36)$$

$$\{F_m, F_n\} = 2L_{m+n} + \frac{C}{3}(m^2 - 1)\delta_{m,-n}, \quad m \neq 0. \quad (37)$$

The central charge C of the action (5) is 26.

In this superstring theory, there are four bosons (C=4), twenty two transverse fermions (C=11) $(\psi_j^{1,2}, \phi_k^{1,2})$ belonging to two transverse superfermionic modes $\Psi^{1,2}$ (equation 9), the usual conformal ghosts [11, 12] (C=-26) and lastly the superconformal ghosts (β, γ) contributing a central charge 11 in a supplemented Hilbert space. The last action is identical to the action of the 22 longitudinal ghost modes $(\psi_j^{0,3}, \phi_k^{0,k})$ or two superfermionic modes $\Psi^{0,3}$. This identity has been proved in the reference [9]. Thus the central charge is zero and there is Lorentz invariance and the superstring theory is anomaly free.

The physical states are defined as

$$\text{NS:} \quad (L_o - 1)|\phi\rangle = 0, \quad L_m|\phi\rangle = 0, \quad G_r|\phi\rangle = 0, \quad r, m > 0. \quad (38)$$

$$\text{R:} \quad (L_o - 1)|\psi\rangle_\alpha = (F_o^2 - 1)|\psi\rangle_\alpha = 0, \quad L_m|\psi\rangle_\alpha = 0, \quad F_r|\psi\rangle_\alpha = 0, \quad m > 0. \quad (39)$$

α is the spinor component index. Admissible Fock space states are

$$\text{NS Eigenstates :} \quad \prod_{n,\mu} \prod_{m,\nu} \{\alpha_{-n}^\mu\} \{B_{-m}^\nu\} |0\rangle, \quad (40)$$

$$\text{R Eigenstates :} \quad \prod_{n,\mu} \prod_{m,\nu} \{\alpha_{-n}^\mu\} \{D_{-m}^\nu\} |0\rangle. \quad (41)$$

G.S.O. projection operators $(1 + (-1)^F)$ [11] is implicit for the NS eigenstates. Supersymmetry forbids the existence of tachyonic vacuum. Here, there are two vacua which are tachyonic in the scalar NS sector and the other in the fermionic Ramond sector. Their self energies cancel as in the supersymmetry, not showing up in overall Fock space.

In the Ramond or Fermionic sector, I aim at the $\frac{3}{2}$ -spin gravitino. In four flat dimensions, the Rarita-Schwinger equation is

$$\epsilon^{\mu\nu\lambda\sigma}\gamma_5\gamma_\lambda\partial_\nu\psi_\sigma = (g^{\mu\nu}\gamma^\sigma - g^{\mu\sigma}\gamma^\nu + g^{\nu\sigma}\gamma^\mu - \gamma^\mu\gamma^\nu\gamma^\sigma)\partial_\nu\psi_\sigma = 0 \quad (42)$$

As shown by GSO in reference [15], this is equivalent to

$$p^\mu\psi_\mu = \gamma^\mu\psi_\mu = \gamma \cdot \psi_\mu = 0 \quad (43)$$

in off-shell momentum space. Condition (43) ensures that there is no admixture of spin $\frac{1}{2}$ fermions.

Let the Fock ground state above the spurious tachyonic state of the R-sector be

$$|\phi_o\rangle = \alpha_{-1}^\mu|0, p\rangle = u_\mu \text{ or } D_{-1}^\mu|0, p\rangle = u_{1\mu}; \quad (44)$$

$u_\mu, u_{1\mu}$ are spinors. This is possible because there is Ramond tachyon along with Neveu-Schwartz. In either case $L_0|\phi_o\rangle = 0$, so that this state is massless. Constraints $L_1|\phi_o\rangle = 0$ and $F_1|\phi_o\rangle = 0$ imply that

$$p^\mu u_\mu = 0, \quad \gamma^\mu u_\mu = 0 \quad (45)$$

It is known that $D_o^\mu \sim \gamma^\mu$, the Dirac gamma matrices and $\alpha_0^\mu = p^\mu$. So $p^\mu u_{1\mu} = 0$ and $\gamma^\mu u_{1\mu} = 0$ as well. There is the freedom to choose a four component spinor ψ^μ such that $u_{1\mu} = \gamma^5 u_\mu$ and

$$\psi_\mu = (1 + \gamma_5)u_\mu \quad (46)$$

By virtue of the above equations

$$\gamma^\mu\psi_\mu = p^\mu\psi_\mu = 0 \quad (47)$$

Further, F_0 is essentially $\gamma \cdot p$. The wave equation is

$$F_0\psi_\mu \sim \gamma \cdot p\psi_\mu = 0 \quad (48)$$

So ψ_μ is the Rarita-Schwinger spin $\frac{3}{2}$ vectorial spinor without spin $\frac{1}{2}$ component. The general relativity equation can be found by invoking the principle of equivalence. The every curved space time point on the manifold, a tangent space is constructed. The vector indices are a, b, .. The vierbein matrices like e_μ^a takes the x space objects to the tangent space and vice versa

$$e_\mu^a e_\nu^b = g^{\mu\nu}, \quad e_\mu^a e^{b\mu} = \delta^{ab} \quad (49)$$

For completely specifying the spin connections ω_μ^{ab} , the covariant derivative D_μ is such that

$$D_\mu e_\lambda^a = 0 \quad (50)$$

With definition of γ matrices

$$\gamma^a e_a^\mu = \gamma^\mu, \quad (51)$$

the covariant derivative of a spinor ψ is

$$D_\mu\psi = (\partial_\mu + \omega_\mu^{ab}\sigma_{ab})\psi \quad (52)$$

σ^{ab} is the antisymmetric product of two gamma matrices. Basically, the principle of equivalence states that in the curved space time, the Rarita-Schwinger equation is

$$\gamma_\nu\gamma^5 D_\sigma\psi_\rho \epsilon^{\mu\nu\sigma\rho} = 0 \quad (53)$$

With $e = \sqrt{g}$, the invariant action in the curved space time is

$$S_{gravitino} = -\frac{i}{2} \int d^4x \, e \, \bar{\psi}_\mu \gamma_\nu \gamma^5 D_\sigma \psi_\rho \, \epsilon^{\mu\nu\sigma\rho} \quad (54)$$

For the graviton, the problem is more involved but interesting. Let C^{ij} be the coefficients of the probabilities of emission and absorption of the superpartners Ψ^μ in the i^{th} and j^{th} sites at the same instance. Then consider the string state(dropping the creation operator suffix '-1/2') above the spurious NS tachyonic state,

$$h^{\dagger\mu\nu}(p) = \sum_{i,j} C^{ij} (b_i^\mu b_j^\nu + b_i^\nu b_j^\mu - 2\eta^{\mu\nu} b_i^\lambda b_j^\lambda) |0, p\rangle. \quad (55)$$

The presence of NS tachyon help build the massless graviton state. This is symmetric and traceless. Further, L_0 will be taken as the free Hamiltonian H_o in the interaction representation. Operating on this state, one gets $L_0 h^{\dagger\mu\nu}(p) = 0$. So this is massless. Also $p_\mu h^{\dagger\mu\nu}(p) = p_\nu h^{\dagger\mu\nu}(p) = 0$ if $C^{ij} = C^{ji}$. If C^{ij} is symmetric, a little algebra shows that

$$G_{\frac{1}{2}} h_{\mu\nu}^\dagger(p) = \left[G_{\frac{1}{2}}, h_{\mu\nu}^\dagger(p) \right] = 0. \quad (56)$$

Thus $h_{\mu\nu}^\dagger(p)$ satisfy all physical state conditions due to Virasoro algebraic relations. This should be graviton. The commutator is

$$[h_{\mu\nu}(p), h_{\mu\nu}(q)] = 2\pi |C|^2 \delta^4(p - q). \quad (57)$$

To switch over quantum field theory, we define the gravitational field by space time Fourier transform of $h_{\mu\nu}(p)$

$$h_{\mu\nu}(x) = \frac{1}{(2\pi)^3} \int \frac{d^3 p}{\sqrt{2p_0}} [h_{\mu\nu}(p) e^{ipx} + h_{\mu\nu}^\dagger(p) e^{-ipx}], \quad (58)$$

with the commutator

$$[h_{\mu\nu}(x), h_{\lambda\sigma}(y)] = \frac{1}{(2\pi)^3} \int \frac{d^3 p}{2p_0} [e^{ip \cdot (x-y)} - e^{-ip \cdot (x-y)}] f^{\mu\nu, \lambda\sigma}, \quad (59)$$

where

$$f^{\mu\nu, \lambda\sigma} = g^{\mu\lambda} g^{\nu\sigma} + g^{\nu\lambda} g^{\mu\sigma} - g^{\mu\nu} g^{\sigma\lambda}. \quad (60)$$

The Feynman propagator is

$$\Delta^{\mu\nu, \lambda\sigma}(x - y) = \langle 0 | T h_{\mu\nu}(x) h_{\lambda\sigma}(y) | 0 \rangle = \frac{1}{(2\pi)^4} \int d^4 p \Delta^{\mu\nu, \lambda\sigma}(p) e^{ip \cdot (x-y)}, \quad (61)$$

where

$$\Delta^{\mu\nu, \lambda\sigma}(p) = \frac{1}{2} f^{\mu\nu, \lambda\sigma} \frac{1}{p^2 - i\epsilon}. \quad (62)$$

This is the propagator of the graviton in the interaction representation. But we do not yet have the general relativity. I take the clue from the case of gravitino and follows it up. In four dimensions, the fourier transform is

$$h_{\mu\nu}(x) = \int \frac{d^4 p}{(2\pi)^4} h_{\mu\nu}(p) e^{ipx} \quad (63)$$

and the hermitian

$$h_{\mu\nu}^\dagger(x) = \int \frac{d^4 p}{(2\pi)^4} h_{\mu\nu}^\dagger(p) e^{-ipx} \quad (64)$$

They are related by

$$h_\lambda^\dagger h_\nu^\lambda = \delta_{\mu\nu} \quad (65)$$

by choosing the normalisation

$$\langle 0, p | 0, q \rangle = (2\pi)^4 \delta^4(p - q) \quad (66)$$

and absorbing $\delta^{(4)}(0)$ in $|C|^2$. In tangent space,

$$h_{\mu\nu}(x) = e_\mu^a(x)e_\nu^b(x)T_{ab} \quad (67)$$

with $h_{\mu\mu}(x) = T_{ab}=0$. After some calculation, one gets

$$h_{\mu\nu}(x) = e_\mu^a e_\nu^b(x)T_{ab}, \quad \text{with} \quad h_{\mu\mu}(x) = T_{bb} = 0. \quad (68)$$

After some calculation, one gets by equation(68)

$$[D_\mu, D_\lambda] h_\nu^\lambda = e_a^\lambda e_{\nu b} R_{\mu\lambda}^{ac} T^{cb} = 0 \quad (69)$$

where the Riemannian Curvature tensor is

$$R_{\mu\lambda}^{ac} = \partial_\mu \omega_\lambda^{ac} + \partial_\mu \omega_\lambda^{dc} - (\mu \leftrightarrow \lambda) \quad (70)$$

Inverting equation(69)

$$[D_\mu, D_\lambda] h_\nu^\lambda = R_{\mu\lambda} h_\nu^\lambda = 0 \quad (71)$$

as the parallel transport relation. Using the equation (65), one gets

$$R_{\mu\nu} = h_\nu^{\dagger\lambda} R_{\mu\sigma} h_\lambda^\sigma = h_\nu^{\dagger\lambda} [D_\mu, D_\sigma] h_\lambda^\sigma \quad (72)$$

At this point, it is appropriate to quote the words of Misner, Thorne and Wheeler [16] 'The Laws of Physics written in abstract geometric form, differ in no way whatsoever between curved space time and flat space time; this is guaranteed by and in fact is a mere rewording of the equivalence principle. In equation (72), the right hand side $[D_\mu, D_\sigma] = [\partial_\mu, \partial_\nu] = 0$ in flat space time. So anywhere in the universe with curved or flat space time

$$R_{\mu\nu} = 0 \quad (73)$$

This is the Einstein equation of General Relativity deduced from string theory. The graviton action is, with $R = g^{\mu\nu} R_{\mu\nu}$,

$$S_{gravitation} = -\frac{1}{2\kappa} \int d^4x \, e \, R \quad (74)$$

κ is the gravitational coupling constant, whose square is proportional to the Newton's constant. The N=1 supergravity action in four dimensions is the sum of equations (43) and (74),

$$S = -\frac{1}{2} \int d^4x \, e \left(\frac{1}{\kappa^2} R + \bar{\psi}_\mu \gamma_\nu \gamma^5 D_\sigma \psi_\rho \, \epsilon^{\mu\nu\sigma\rho} \right) \quad (75)$$

The local supergravity transformations that leave it invariant are

$$\delta\psi_\mu = \frac{1}{\kappa} D_\mu \epsilon(x) \quad \text{and} \quad \delta e_\mu^m = -\frac{i}{2} \kappa \bar{\epsilon}(x) \gamma^m \psi_\mu \quad (76)$$

I have succeeded in deriving the 4d, N=1 supergravity theory as it develops from a renormalisable open 26d bosonic string. This has unlimited useful consequences in physics.

I thank Prof L. Maharana but for whose discussions and suggestions, this article could not have been completed.

-
- [1] S. Desser and B. Zumino, Phys. Letts. **62B**, 335 (1976); Phys. Letts. **65B**, 369(1976).
 - [2] Y. Nambu, *Lectures at Copenhagen Symposium* (1970);
 - [3] T. Goto, Prog. Theo. Phys. **46**, 1566 (1971).
 - [4] A. Casher, F. Englert, H. Nicolai and A. Taormina, Phys. Letts. **B162**, 121 (1985).
 - [5] F. Englert, A. Hourt and A. Taormina, JHEP, 0108.013, 1 (2001).
 - [6] B. B. Deo, Phys. Letts. **B557**, 115 (2003) and "Supersymmetric Standard Model from String Theory, ArXive hep-th/03010441;

- [7] B. B. Deo and L. Maharana, Mod. Phys. Letts. **19A**, 1 (2004).
- [8] B. B. Deo and L. Maharana, *Derivation of Einstein Equation from a Four Dimensionai Superstring*, hep-th/0212004 (to be published in Cosmology and Gravitation).
- [9] B. B. Deo and L. Maharana, 'A Novel string in four dimensions from bosonic String with some Application', ArXive hep-th/0307172 (to be published in Int. Jour. Mod. Phys. A).
- [10] S. Mandelstam, Phys. Rev. **D11**, 3026(1975)
- [11] M. B. Green, J. H. Schwartz and E. Witten, *Superstring Theory*, Vol.I, Cambridge University Press, England (1987).
- [12] M. Kaku, *Introduction to Superstring and M. Theory*, 2nd Edn, Springer, New York (1999).
- [13] A. Neveu and J. H. Schwarz, Nucl. Phys. **B31**, 86(1971).
- [14] P. Ramond, Phys. Rev. **D3**, 2415(1971).
- [15] F. Gliozzi, J. Scherk and D. Olive, Nucl. Phys. **B122**, 253 (1977).
- [16] C. W. Misner, K. S. Thorne, J. A. Wheeler, *Gravitation*, W. H. Freeman and Company, Sanfrancisco, 1970.